

The generalized Markov measure as an equilibrium state.

Ivan Werner*

Email: ivan_werner@pochta.ru

February 1, 2008

Abstract

In this paper, we continue development of the theory of contractive Markov systems (CMS) initiated in [13]. Also, this work can be seen as a small contribution to the theory of equilibrium states.

We construct an energy function on the code space, using the coding map from [15], and show that the generalized Markov measure associated with an irreducible CMS is a unique equilibrium state for this energy function if the vertex sets form an open partition of the state space of the CMS and the restrictions of the probability functions on their vertex sets are Dini-continuous and bounded away from zero.

MSC: 37D35, 28D05, 28A80, 37H99, 60J05.

Keywords: equilibrium states, contractive Markov systems (CMS), iterated function systems (IFS) with place-dependent probabilities, g -measures, random systems with complete connections, Markov chains, fractals.

1 Introduction

In [13], we introduced a theory of *contractive Markov systems (CMS)* which provides a unifying framework in so-called 'fractal' geometry. It extends the known theory of *iterated function systems (IFS) with place dependent probabilities*, which are contractive on average, [1][3] in a way that it also covers *graph directed constructions* of 'fractal' sets [8]. In particular, Markov chains associated with such systems naturally extend finite Markov chains and inherit some of their properties.

*The work was partially supported by EPSRC and School of Mathematics and Statistics of University of St Andrews.

By a *Markov system* we mean the following structure on a metric space (K, d) , which generates a Markov process. It is given by a family

$$(K_{i(e)}, w_e, p_e)_{e \in E}$$

(see Fig. 1) where E is the set of edges of a finite directed (multi)graph (V, E, i, t) ($V := \{1, \dots, N\}$ is the set of vertices of the directed (multi)graph (we do not exclude the case $N = 1$), $i : E \rightarrow V$ is a map indicating the initial vertex of each edge and $t : E \rightarrow V$ is a map indicating the terminal vertex of each edge), $K_1, K_2, \dots, K_N$ is a partition of the metric space K into non-empty Borel subsets, $(w_e)_{e \in E}$ is a family of Borel measurable self-maps on the metric space such that $w_e(K_{i(e)}) \subset K_{t(e)}$ for all $e \in E$ and $(p_e)_{e \in E}$ is a family of Borel measurable probability functions on K (i.e. $p_e(x) \geq 0$ for all $e \in E$ and $\sum_{e \in E} p_e(x) = 1$ for all $x \in K$) (associated with the maps) such that each p_e is zero on the complement of $K_{i(e)}$.




Fig. 1. A Markov system

A Markov system is called *irreducible* or *aperiodic* iff its directed graph is irreducible or aperiodic respectively. We call a Markov system $(K_{i(e)}, w_e, p_e)_{e \in E}$ *contractive* with an *average contracting rate* $0 < a < 1$ iff it satisfies the following *condition of contractiveness on average*:

$$\sum_{e \in E} p_e(x) d(w_e x, w_e y) \leq a d(x, y) \text{ for all } x, y \in K_i, i = 1, \dots, N. \quad (1)$$

This condition was discovered by R. Isaac in 1961 [4] (for the case $N = 1$).

Markov system $(K_{i(e)}, w_e, p_e)_{e \in E}$ determines a Markov operator U on the set of all bounded Borel measurable functions $\mathcal{L}^0(K)$ by

$$Uf := \sum_{e \in E} p_e f \circ w_e \text{ for all } f \in \mathcal{L}^0(K)$$

and its adjoint operator U^* on the set of all Borel probability measures $P(K)$ by

$$U^*\nu(f) := \int U(f)d\nu \text{ for all } f \in \mathcal{L}^0(K) \text{ and } \nu \in P(K).$$

Remark 1 Note that each map w_e and each probability p_e need to be defined only on the corresponding vertex set $K_{i(e)}$. This is sufficient for the condition (1) and the definition of U^* . For the definition of U , we can consider each w_e to be extended on the whole space K arbitrarily and each p_e to be extended on K by zero.

Also, the situation applies where each vertex set K_i has its own metric d_i . In this case, one can set

$$d(x, y) = \begin{cases} d_i(x, y) & \text{if } x, y \in K_i \\ \infty & \text{otherwise} \end{cases}$$

and use the convention $0 \times \infty = 0$.

We say $\mu \in P(K)$ is an invariant measure of the CMS iff $U^*\mu = \mu$. A Borel probability measure μ is called *attractive* measure of the CMS if

$$U^{*n}\nu \xrightarrow{w^*} \mu \text{ for all } \nu \in P(K),$$

where w^* means weak* convergence. Note that an attractive probability measure is a unique invariant probability measure of the CMS if U maps continuous functions on continuous functions. The following theorem states some properties of such CMSs.

Theorem 1 *Let K be a metric space in which sets of finite diameter are relatively compact. Suppose $(K_{i(e)}, w_e, p_e)_{e \in E}$ is a CMS with an average contracting rate $0 < a < 1$ such that the family $K_1, \dots, K_N$ partitions K into open subsets and each $p_e|_{K_{i(e)}}$ is continuous on $K_{i(e)}$. Then the following hold:*

- (i) *The sequence $(U^{*k}\delta_x)_{k \in \mathbb{N}}$ is tight for all $x \in K$, i.e. for all $\epsilon > 0$, there exists a compact subset $Q \subset K$ such that $U^{*k}\delta_x(Q) \geq 1 - \epsilon$ for all $k \in \mathbb{N}$.*
- (ii) *The CMS has an invariant Borel probability measure μ .*
- (iii) *The invariant probability measure μ is unique iff*

$$\frac{1}{n} \sum_{k=1}^n U^k g(x) \rightarrow \int g d\mu \text{ for all } x \in K \text{ and } g \in C_B(K).$$

- (iv) *If the invariant probability measure is unique, then*

$$\sum_{i=1}^N \int_{K_i} d(x, x_i) d\mu(x) < \infty \text{ for all } x_i \in K_i, i = 1, \dots, N.$$

For the proof see [13].

Furthermore, it was shown in [13] that contractive Markov systems inherit some properties of finite Markov chains if the restrictions of the probabilities on their vertex sets satisfy some stronger conditions. Namely, it has a unique invariant probability measure in an irreducible case and an attractive probability measure in an aperiodic case if the vertex sets $K_1, \dots, K_N$ form an open partition of the state space and each $p_e|_{K_{i(e)}}$ is Dini-continuous and bounded away from zero (see also [14] for the case of constant probabilities $p_e|_{K_{i(e)}}$ and compact state space).

A function $h : (X, d) \longrightarrow \mathbb{R}$ is called *Dini-continuous* iff for some $c > 0$

$$\int_0^c \frac{\phi(t)}{t} dt < \infty$$

where ϕ is the modulus of uniform continuity of h , i.e.

$$\phi(t) := \sup\{|h(x) - h(y)| : d(x, y) \leq t, x, y \in X\}.$$

It is easily seen that Dini-continuity is weaker than Hölder and stronger than the uniform continuity. There is a well known characterization of Dini-continuity, which will be useful later.

Lemma 1 *Let $0 < c < 1$ and $b > 0$. A function h is Dini-continuous iff*

$$\sum_{n=0}^{\infty} \phi(bc^n) < \infty$$

where ϕ is the modulus of uniform continuity of h .

The proof is simple (e.g. see [13]).

Also, associated with the Markov system is a measure preserving transformation $S : (\Sigma, \mathcal{B}(\Sigma), M) \longrightarrow (\Sigma, \mathcal{B}(\Sigma), M)$, which we call a *generalized Markov shift*, where $\Sigma := \{(\dots, \sigma_{-1}, \sigma_0, \sigma_1, \dots) : \sigma_i \in E \forall i \in \mathbb{Z}\}$ is the *code space* provided with the product topology, $\mathcal{B}(\Sigma)$ denotes Borel σ -algebra on Σ and M is a *generalized Markov measure* on $\mathcal{B}(\Sigma)$ given by

$$M(m[e_1, \dots, e_k]) := \int p_{e_1}(x) p_{e_2}(w_{e_1}x) \dots p_{e_k}(w_{e_{k-1}} \circ \dots \circ w_{e_1}x) d\mu(x)$$

for every cylinder set $m[e_1, \dots, e_k] := \{\sigma \in \Sigma : \sigma_m = e_1, \dots, \sigma_{m+k-1} = e_k\}$, $m \in \mathbb{Z}$, where μ is an invariant Borel probability measure of the Markov system, and S is the usual left shift map on Σ . It is easy to verify that S preserves measure M , since $U^*\mu = \mu$ (see [15]).

For a contractive Markov system (CMS), the Markov process associated with the CMS can be represented as a factor of the generalized Markov shift via a

coding map $F : (\Sigma, \mathcal{B}(\Sigma), M) \longrightarrow K$ which was constructed in [15]. It is defined by

$$F_{x_1, \dots, x_N}(\sigma) := \lim_{m \rightarrow -\infty} w_{\sigma_0} \circ w_{\sigma_{-1}} \circ \dots \circ w_{\sigma_m} x_{i(\sigma_m)} \text{ for } M\text{-a.e. } \sigma \in \Sigma,$$

under some conditions (see the next lemma), where $x_i \in K_i$ for each $i = 1, \dots, N$.

Lemma 2 *Suppose that $p_e|_{K_{i(e)}}$ is Dini-continuous and there exists $\delta > 0$ such that $p_e|_{K_{i(e)}} \geq \delta$ for all $e \in E$. Let $x_i, y_i \in K_i$ for all $1 \leq i \leq N$. Then the following hold:*

- (i) $F_{x_1 \dots x_N}$ is defined M -a.e.,
- (ii) $F_{x_1 \dots x_N} = F_{y_1 \dots y_N}$ M -a.e. and
- (iii) There exists a sequence of closed subsets $Q_1 \subset Q_2 \subset \dots \subset \Sigma$ with $\lim_{k \rightarrow \infty} M(Q_k) = 1$ such that all $F_{x_1 \dots x_N}|_{Q_k}$ are locally Hölder-continuous with the same Hölder-constants.

In the following, we fix $x_1, \dots, x_N$ and denote the coding map simply by F . This coding map is the key tool for our investigation.

Example 1 (decimal expansion) Consider ten maps $w_e, e \in E := \{0, \dots, 9\}$, on $([0, 1], |\cdot|)$ given by $w_e(x) := 1/10x + e/10$ for all $x \in [0, 1]$. Obviously, for any family of probability functions $p_e, e \in E$, the family $([0, 1], w_e, p_e)_{e \in E}$ is a CMS. The coding map for this CMS is nothing else as the usual decimal expansion of real numbers from $[0, 1]$.

Example 2 Let $G := (V, E, i, t)$ be a finite irreducible directed (multi)graph. Let $\Sigma_G^- := \{(\dots, \sigma_{-1}, \sigma_0) : \sigma_m \in E \text{ and } t(\sigma_m) = i(\sigma_{m-1}) \forall m \in \mathbb{Z} \setminus \mathbb{N}\}$ (one-sided subshift of finite type associated with G) endowed with the metric $d(\sigma, \sigma') := 2^k$ where k is the smallest integer with $\sigma_i = \sigma'_i$ for all $k < i \leq 0$. Let g be a positive, Dini-continuous function on Σ_G such that

$$\sum_{y \in T^{-1}(\{x\})} g(y) = 1 \text{ for all } x \in \Sigma_G$$

where T is the right shift map on Σ_G^- . Set $K_i := \{\sigma \in \Sigma_G^- : t(\sigma_0) = i\}$ for every $i \in V$ and, for every $e \in E$,

$$w_e(\sigma) := (\dots, \sigma_{-1}, \sigma_0, e), \quad p_e(\sigma) := g(\dots, \sigma_{-1}, \sigma_0, e) \text{ for all } \sigma \in K_{i(e)}.$$

Obviously, maps $(w_e)_{e \in E}$ are contractions. Therefore, $(K_{i(e)}, w_e, p_e)_{e \in E}$ defines a CMS. An invariant probability measure of such a CMS is called a g -measure. This notion was introduced by M. Keane [5]. In this example, the coding map $F : \Sigma_G \longrightarrow \Sigma_G^-$ is nothing else as the natural projection, and the generalized Markov measure is the natural extension of the g -measure (or in other words, the g -measure is the image of the generalized Markov measure under the coding

map). The equivalence between g -measures and equilibrium states for $\log g$ was elaborated by Ledrappier [7]. See also [11] for a proof of the uniqueness of the equilibrium state in this example.

In this paper, we are concerned with the following question. Is the generalized Markov measure associated with a CMS a unique equilibrium state for some energy function? We report here that the answer to this question is YES, under some conditions (see Corollary 1). The result seems to be beyond the well known thermodynamic formalism. It turns out that our energy function is not upper semicontinuous in general (see Example 3). Recall that the upper semicontinuity of an energy function is a widely used condition in the rigorous theory of equilibrium states which insures that the convex set of equilibrium states is non-empty and compact in the weak* topology (see e.g. [6]).

Also, an interesting point of the presented result is that it introduces some symbolic dynamical systems of infinite type, which have the full measures M .

Definition 1 Let X be a metric space and T a continuous transformation on it. Denote by $P(X)$ the set of all Borel probability measures on X and by $P_T(X)$ the set of all T -invariant Borel probability measures on X . We call a Borel measurable function $f : X \rightarrow [-\infty, 0]$ an *energy function*. Suppose that T has a finite topological entropy, i.e. $\sup_{\Theta \in P_T(X)} h_\Theta(T) < \infty$, where $h_\Theta(T)$ is the Kolmogorov-Sinai entropy of T with respect to measure Θ . We call

$$P(f) = \sup_{\Theta \in P_T(X)} (h_\Theta(T) + \Theta(f))$$

the *pressure* of f . We call $\Lambda \in P_T(X)$ an *equilibrium state* for f iff

$$h_\Lambda(T) + \Lambda(f) = P(f).$$

Note that $\sup_{\Theta \in P_S(\Sigma)} h_\Theta(S) = \log |E|$ (e.g. Example 4.2.6 in [6]).

2 Main part

Let

$$\mathcal{M} := (K_{i(e)}, w_e, p_e)_{e \in E}$$

be a contractive Markov system with an average contracting rate $0 < a < 1$ and an invariant Borel probability measure μ . We assume that: (K, d) is a metric space in which sets of finite diameter are relatively compact and the family $K_1, \dots, K_N$ partitions K into non-empty open subsets; each probability function $p_e|_{K_{i(e)}}$ is uniformly continuous and bounded away from zero by $\delta > 0$; the set of edges E is finite and the map $i : E \rightarrow V$ is surjective. Note that the assumption on the metric space implies that it is locally compact separable and

complete. We shall denote the space of all bounded continuous functions on K by $C_B(K)$.

We consider Σ endowed with the metric $d'(\sigma, \sigma') := (1/2)^k$ where k is the largest integer with $\sigma_i = \sigma'_i$ for all $|i| < k$. Fix $x_i \in K_i$ for all $i = 1, \dots, N$.

The construction of the energy function goes through a definition of an appropriate shift invariant subset of Σ on which the energy function shall be finite.

Let

$$\Sigma_G := \{\sigma \in \Sigma : t(\sigma_j) = i(\sigma_{j+1}) \ \forall j \in \mathbb{Z}\},$$

$$D := \{\sigma \in \Sigma_G : \lim_{m \rightarrow -\infty} w_{\sigma_0} \circ w_{\sigma_{-1}} \circ \dots \circ w_{\sigma_m} x_{i(\sigma_m)} \text{ exists}\}$$

and

$$Y := \bigcap_{i=-\infty}^{\infty} S^i(D).$$

Lemma 3 (i) $F(\sigma)$ is defined for all $\sigma \in Y$ and Y is a shift invariant subset of Σ_G .

(ii) If each $p_e|_{K_{i(e)}}$ is Dini-continuous, then $M(Y) = 1$.

(iii) If each $p_e|_{K_{i(e)}}$ is Dini-continuous and the CMS has an invariant probability measure μ such that $\mu(K_{i(e)}) > 0$ for all $e \in E$. Then Y is dense in Σ_G .

Proof. (i) is clear, by the definitions of F and Y . Note that the condition of the contractiveness on average and the boundedness away from zero of the functions $p_e|_{K_{i(e)}}$ imply that each map $w_e|_{K_{i(e)}}$ is continuous (Lipschitz). Therefore, $S(D) \subset D$. By Lemma 2, $M(D) = 1$. This implies that $M(Y) = 1$. If in addition $\mu(K_{i(e)}) > 0$ for all $e \in E$, then $M(O) > 0$ for every open $O \subset \Sigma_G$. This implies (iii). $\square$

Remark 2 (i) Note that Y and F depend on the choice of x_i 's. By Corollary 1 in [15], Y changes only modulo M -zero set by a different choice of x_i 's.

(ii) If all maps $w_e|_{K_{i(e)}}$ are contractive, then $Y = \Sigma_G$ and $F|_{\Sigma_G}$ is Hölder-continuous (easy to check).

In the following, we are going to represent the generalized Markov measure M as a unique equilibrium state for the energy function u given by

$$u(\sigma) = \begin{cases} \log p_{\sigma_1} \circ F(\sigma) & \text{if } \sigma \in Y \\ -\infty & \text{if } \sigma \in \Sigma \setminus Y \end{cases} \quad (2)$$

if the CMS has a unique invariant Borel probability measure and $M(Y) = 1$.

Let's consider a simple example which illustrates F , Y and u .

Example 3 Let $(K, d) = (\mathbb{R}, |\cdot|)$. Consider two maps

$$w_0(x) := \frac{1}{2}x, \quad w_1(x) := 2x \text{ for all } x \in \mathbb{R}$$

with probability functions

$$p_0(x) := \frac{1}{6} \sin^2 x + \frac{17}{24}, \quad p_1(x) := \frac{1}{6} \cos^2 x + \frac{1}{8} \text{ for all } x \in \mathbb{R}.$$

Then a simple calculation shows that $(\mathbb{R}, w_e, p_e)_{e \in \{0,1\}}$ defines a CMS with an average contracting rate $45/48$. In this case, $\Sigma_G = \{0,1\}^{\mathbb{Z}}$. If we take $x = 0$ for the definition of Y , then, obviously, $Y = \Sigma_G$. Now, let $x \neq 0$. Let $N_{0n}(\sigma)$ and $N_{1n}(\sigma)$ be the numbers of zeros and ones in $(\sigma_{-n}, \dots, \sigma_0)$ respectively for every $\sigma \in \Sigma_G$. Then, obviously, $\sigma \notin Y$ if $(N_{1n}(\sigma) - N_{0n}(\sigma)) \rightarrow \infty$. Hence, $Y \neq \Sigma_G$ and, by Lemma 3 (iii), Y is a dense shift invariant subset of Σ_G . Since Y is not closed, u is not upper semicontinuous.

Also, it is not difficult to see that in a general case there is no hope to find x_i such that u becomes upper semicontinuous, e.g. change w_1 to $w_1(x) = 2x + 1$, then, for any choice of x for the definition of Y , $Y \neq \Sigma_G$.

In what follows, we shall denote the restrictions of F and M to Y by the same letters.

Definition 2 Let $\mathcal{A}_0$ be the finite σ -algebra on Σ generated by the zero-time partition. Set $\mathcal{F} := \bigvee_{i=0}^{\infty} S^i \mathcal{A}_0$. Let $\Lambda \in P_S(\Sigma)$. Define, for $e \in E$,

$$g_{\Lambda e} := E_{\Lambda}(1_{1[e]}|\mathcal{F}) \text{ and } g_{\Lambda}(\sigma) := g_{\Lambda \sigma_1}(\sigma) \text{ for } \Lambda\text{-a.e. } \sigma \in \Sigma,$$

where $E_{\Lambda}(\cdot|\cdot)$ denotes the conditional expectation with respect to Λ . Note that we can consider $g_{\Lambda e}(\sigma) = g_{\Lambda e}(\dots, \sigma_{-1}, \sigma_0)$ for all $\sigma \in \Sigma$.

We are going to show that every $\Lambda \in P_S(\Sigma)$ is an equilibrium state for a version of $\log g_{\Lambda}$, but first let's see some properties of the function g_{Λ} .

Lemma 4 Let $\Lambda \in P_S(\Sigma)$.

- (i) $0 \leq g_{\Lambda} \leq 1$ Λ -a.e..
- (ii) $\sum_{e \in E} g_{\Lambda e} = 1$ Λ -a.e..
- (iii) $\Lambda(\{g_{\Lambda} = 0\}) = 0$.
- (iv) $1/g_{\Lambda}, \log g_{\Lambda} \in \mathcal{L}^1(\Lambda)$.

Proof. (i) and (ii) are clear by the properties of the conditional expectation. For (iii), observe that

$$\begin{aligned} \Lambda(\{g_{\Lambda} = 0\}) &= \sum_{e \in E_{\{g_{\Lambda}=0\}}} \int 1_{1[e]} d\Lambda \\ &= \sum_{e \in E_{\{g_{\Lambda e}=0\}}} \int g_{\Lambda e} d\Lambda = 0. \end{aligned}$$

For (iv), by the pull-out property of the conditional expectation (see e.g. Theorem 10.1.9 in [2]),

$$\begin{aligned}
\int \left(\frac{1}{g_\Lambda} \wedge n \right) d\Lambda &= \sum_{e \in E} \int 1_{1[e]} \left(\frac{1}{g_{\Lambda e}} \wedge n \right) d\Lambda \\
&= \sum_{e \in E} \int E_\Lambda \left(1_{1[e]} \left(\frac{1}{g_{\Lambda e}} \wedge n \right) \middle| \mathcal{F} \right) d\Lambda \\
&= \sum_{e \in E} \int g_{\Lambda e} \left(\frac{1}{g_{\Lambda e}} \wedge n \right) d\Lambda \\
&\leq \sum_{e \in E} 1,
\end{aligned}$$

for all $n \in \mathbb{N}$. Hence, $1/g_\Lambda \in \mathcal{L}^1(\Lambda)$. Since $\log x \leq x - 1$,

$$\begin{aligned}
\int |\log g_\Lambda| d\Lambda &= \int \log \frac{1}{g_\Lambda} d\Lambda \\
&\leq \int \left(\frac{1}{g_\Lambda} - 1 \right) d\Lambda \\
&< \infty.
\end{aligned}$$

Thus, $\log g_\Lambda \in \mathcal{L}^1(\Lambda)$. □

Lemma 5 *Let $\Lambda \in P_S(\Sigma)$. For each $e \in E$, let $\dot{g}_{\Lambda e}$ be a Borel measurable version of $g_{\Lambda e}$ such that $\log \dot{g}_\Lambda$ is an energy function and $\sum_{e \in E} \dot{g}_{\Lambda e}(\sigma) \leq 1$ for all $\sigma \in \Sigma$. Then Λ is an equilibrium state for $\log \dot{g}_\Lambda$ and*

$$h_\Lambda(S) + \Lambda(\log \dot{g}_\Lambda) = 0,$$

i.e. $P(\log \dot{g}_\Lambda) = 0$. If $\Theta \in P_S(\Sigma)$ is also an equilibrium state for $\log \dot{g}_\Lambda$, then $g_\Theta = \dot{g}_\Lambda$ Θ -a.e..

Proof. The proof goes similarly to that of Theorem 1 in [7]. The key for the proof is the pull-out property of the conditional expectation (see e.g. Theorem 10.1.9 in [2]).

Since $\mathcal{A}_0$ is a generator for the Borel σ -algebra, we know (see e.g. [12]) that

$$h_\Lambda(S) = - \sum_{e \in E} \int E_\Lambda \left(1_{1[e]} \middle| \mathcal{F} \right) \log E_\Lambda \left(1_{1[e]} \middle| \mathcal{F} \right) d\Lambda.$$

By Lemma 2, $1_{1[e]} \log g_{\Lambda e} \in \mathcal{L}^1(\Lambda)$ for all $e \in E$. Therefore, by the pull-out

property of the conditional expectation,

$$\begin{aligned}
h_\Lambda(S) &= - \sum_{e \in E} \int 1_{1[e]} \log g_{\Lambda e} d\Lambda \\
&= - \sum_{e \in E} \int 1_{1[e]} \log g_\Lambda d\Lambda \\
&= - \int \log g_\Lambda d\Lambda.
\end{aligned}$$

For the first part of the Lemma, it remains to show that

$$h_\Theta(S) + \Theta(\log \dot{g}_\Lambda) \leq 0 \text{ for all } \Theta \in P_S(\Sigma).$$

Let $\Theta \in P_S(\Sigma)$. By the above, $h_\Theta(S) = -\Theta(\log g_\Theta)$. If $\Theta(\{\dot{g}_\Lambda = 0\}) > 0$, then $h_\Theta(S) + \Theta(\log \dot{g}_\Lambda) = -\infty < 0$. Otherwise, since $\log x \leq x - 1$ for all $x > 0$, it follows that

$$\begin{aligned}
h_\Theta(S) + \Theta(\log \dot{g}_\Lambda) &= \int \log \frac{\dot{g}_\Lambda}{g_\Theta} d\Theta \\
&\leq \int \left(\frac{\dot{g}_\Lambda}{g_\Theta} - 1 \right) d\Theta.
\end{aligned}$$

By Lemma 2, $1_{1[e]}(\dot{g}_{\Lambda e}/g_{\Theta e} - 1) \in \mathcal{L}^1(\Theta)$ for all $e \in E$. Therefore, by the pull-out property of the conditional expectation,

$$\begin{aligned}
\int \left(\frac{\dot{g}_\Lambda}{g_\Theta} - 1 \right) d\Theta &= \sum_{e \in E} \int 1_{1[e]} \left(\frac{\dot{g}_{\Lambda e}}{g_{\Theta e}} - 1 \right) d\Theta \\
&= \sum_{e \in E} \int g_{\Theta e} \left(\frac{\dot{g}_{\Lambda e}}{g_{\Theta e}} - 1 \right) d\Theta \\
&= \int \sum_{e \in E} (\dot{g}_{\Lambda e} - g_{\Theta e}) d\Theta \\
&\leq 0.
\end{aligned}$$

Hence,

$$h_\Theta(S) + \Theta(\log \dot{g}_\Lambda) \leq 0,$$

i.e. Λ is an equilibrium state for $\log \dot{g}_\Lambda$.

Now, suppose Θ_0 is also an equilibrium state for $\log \dot{g}_\Lambda$, i.e.

$$h_{\Theta_0}(S) + \Theta_0(\log \dot{g}_\Lambda) = \sup_{\Theta \in P_S(\Sigma)} (h_\Theta(S) + \Theta(\log \dot{g}_\Lambda)) = h_\Lambda(S) + \Lambda(\log \dot{g}_\Lambda) = 0.$$

Then, by the above, the following equality must hold true:

$$\int \log \frac{\dot{g}_\Lambda}{g_{\Theta_0}} d\Theta_0 = \int \left(\frac{\dot{g}_\Lambda}{g_{\Theta_0}} - 1 \right) d\Theta_0.$$

But it is true if and only if

$$\log \frac{\dot{g}_\Lambda}{g_{\Theta_0}} = \left(\frac{\dot{g}_\Lambda}{g_{\Theta_0}} - 1 \right) \Theta_0\text{-a.e.}$$

And the latter happens if and only if $\dot{g}_\Lambda = g_{\Theta_0}$ Θ_0 -a.e.. $\square$

Now, we are going to prove what seems to be the main lemma for the generalized Markov shift associated with a contractive Markov system. For that we need to define some measures on the product space $K \times \Sigma$.

Denote by $\mathcal{A}$ the finite σ -algebra generated by the partition $\{0[e] : e \in E\}$ of Σ and define, for each integer $m \leq 1$,

$$\mathcal{A}_m := \bigvee_{i=m}^{+\infty} S^{-i}\mathcal{A},$$

which is the smallest σ -algebra containing all finite σ -algebras $\bigvee_{i=m}^n S^{-i}\mathcal{A}$, $n \geq m$. Let $x \in K$. For every integer $m \leq 1$, let P_x^m be a probability measure on σ -algebra $\mathcal{A}_m$ given by

$$P_x^m(m[e_m, \dots, e_n]) = p_{e_m}(x)p_{e_{m+1}}(w_{e_m}(x)) \dots p_{e_n}(w_{e_{n-1}} \circ \dots \circ w_{e_m}(x))$$

for all cylinders $m[e_m, \dots, e_n]$, $n \geq m$. By Lemma 1 from [15], $x \mapsto P_x^m(A)$ is a Borel measurable function on K . Therefore, we can define, for every integer $m \leq 0$,

$$\tilde{M}_m(A \times Q) := \int_A P_x^m(Q) d\mu(x)$$

for all $A \in \mathcal{B}(K)$ and all $Q \in \mathcal{A}_m$. Then $\tilde{M}_m$ extends uniquely to a probability measure on the product σ -algebra $\mathcal{B}(K) \otimes \mathcal{A}_m$ with

$$\tilde{M}_m(\Omega) = \int P_x^m(\{\sigma \in \Sigma : (x, \sigma) \in \Omega\}) d\mu(x)$$

for all $\Omega \in \mathcal{B}(K) \otimes \mathcal{A}_m$. Note that the set of all $\Omega \in \mathcal{B}(K) \otimes \mathcal{A}_m$ for which the integrand in the above is measurable forms a Dynkin system which contains the set all rectangles $A \times Q$, $A \in \mathcal{B}(K)$, $Q \in \mathcal{A}_m$. As the latter is $\cap$ -stable and generates $\mathcal{B}(K) \otimes \mathcal{A}_m$, the integrand is measurable for all $\Omega \in \mathcal{B}(K) \otimes \mathcal{A}_m$. Further, note that $P_x^m(\{\sigma \in \Sigma : (x, \sigma) \in \Omega\}) = \int 1_\Omega(x, \sigma) dP_x^m(\sigma)$ for all $\Omega \in \mathcal{B}(K) \otimes \mathcal{A}_m$. Therefore

$$\int s d\tilde{M}_m = \int \int s(x, \sigma) dP_x^m(\sigma) d\mu(x)$$

for all $\mathcal{B}(K) \otimes \mathcal{A}_m$ -simple functions s . Now, let ψ be a $\mathcal{B}(K) \otimes \mathcal{A}_m$ -measurable and $\tilde{M}_m$ -integrable function on $K \times \Sigma$. Then the usual monotone approximation of positive and negative parts of ψ by simple functions and the B. Levi Theorem imply that

$$\int \psi d\tilde{M}_m = \int \int \psi(x, \sigma) dP_x^m(\sigma) d\mu(x).$$

Lemma 6 *Suppose $\mathcal{M}$ is a CMS with an invariant Borel probability measure μ such that $C := \sum_{i=1}^N \int_{K_i} d(x, x_i) d\mu(x) < \infty$ for some $x_i \in K$, $i = 1, \dots, N$, and $M(Y) = 1$, where M is the generalized Markov measure associated with $\mathcal{M}$ and μ . Then*

$$E_M(1_{[e]}|\mathcal{F}) = p_e \circ F \text{ } M\text{-a.e. for all } e \in E.$$

Proof. Fix $e \in E$. Let $\mathcal{F}_m := \bigvee_{i=0}^m S^i(\mathcal{A})$ for all $m \in \mathbb{Z} \setminus \mathbb{N}$. Let's use the notation $(\sigma_m, \dots, \sigma_0)^*$ iff $M(m[\sigma_m, \dots, \sigma_0]) > 0$. Then obviously

$$E_M(1_{[e]}|\mathcal{F}_m)(\tilde{\sigma}) = \sum_{(\sigma_m, \dots, \sigma_0)^*} \frac{\int 1_{[e]} dM}{M(m[\sigma_m, \dots, \sigma_0])} 1_{m[\sigma_m, \dots, \sigma_0]}(\tilde{\sigma})$$

for all M -a.e. $\tilde{\sigma} \in \Sigma$. By Doob's Martingale Theorem,

$$E_M(1_{[e]}|\mathcal{F}_m) \rightarrow E_M(1_{[e]}|\mathcal{F}) \text{ } M\text{-a.e..}$$

Now, set

$$Z_m^x(\sigma) := w_{\sigma_0} \circ \dots \circ w_{\sigma_m}(x) \text{ and } Y_m(\sigma) := w_{\sigma_0} \circ \dots \circ w_{\sigma_m}(x_{i(\sigma_m)})$$

for all $x \in K$, $\sigma \in \Sigma$ and $m \leq 0$. Then

$$\begin{aligned} & |E_M(1_{[e]}|\mathcal{F}_m) - p_e \circ F| \\ & \leq \sum_{(\sigma_m, \dots, \sigma_0)^*} \left| \frac{\int_{K \times_m [\sigma_m, \dots, \sigma_0]} p_e \circ Z_m^x(\tilde{\sigma}) d\tilde{M}_m(x, \tilde{\sigma})}{\tilde{M}_m(K \times_m [\sigma_m, \dots, \sigma_0])} - p_e \circ Y_m \right| 1_{m[\sigma_m, \dots, \sigma_0]} + |p_e \circ Y_m - p_e \circ F| \\ & \leq \sum_{(\sigma_m, \dots, \sigma_0)^*} \frac{\int_{K \times_m [\sigma_m, \dots, \sigma_0]} |p_e \circ Z_m^x(\tilde{\sigma}) - p_e \circ Y_m| d\tilde{M}_m(x, \tilde{\sigma})}{\tilde{M}_m(K \times_m [\sigma_m, \dots, \sigma_0])} 1_{m[\sigma_m, \dots, \sigma_0]} + |p_e \circ Y_m - p_e \circ F|. \end{aligned} \tag{3}$$

Set

$$Q_m := \left\{ (x, \sigma) \in K \times \Sigma : d(Z_m^x(\sigma), Y_m(\sigma)) > a^{\frac{-m+1}{2}} C \right\}.$$

Observe that, by the contractiveness on average condition,

$$\begin{aligned} & \int d(Z_m^x(\sigma), Y_m(\sigma)) d\tilde{M}_m(x, \sigma) \\ & = \int \sum_{\sigma_m, \dots, \sigma_0} p_{\sigma_m}(x) \dots p_{\sigma_0}(w_{\sigma_{-1}} \circ \dots \circ w_{\sigma_m} x) d(w_{\sigma_0} \circ \dots \circ w_{\sigma_m} x, w_{\sigma_0} \circ \dots \circ w_{\sigma_m} x_{i(\sigma_m)}) d\mu(x) \\ & \leq a^{-m+1} \sum_{i=1}^N \int_{K_i} d(x, x_i) d\mu(x) \\ & = a^{-m+1} C. \end{aligned}$$

Hence,

$$a^{\frac{-m+1}{2}} C \tilde{M}_m(Q_m) \leq a^{-m+1} C,$$

that is,

$$\tilde{M}_m(Q_m) \leq a^{\frac{-m+1}{2}}.$$

Therefore,

$$(3) \leq X_m + \phi\left(a^{\frac{-m+1}{2}} C\right) + |p_e \circ Y_m - p_e \circ F| \quad M\text{-a.e.}, \quad (4)$$

where

$$X_m := \sum_{(e_m, \dots, e_0)^*} \frac{\tilde{M}_m(Q_m \cap (K \times_m [e_m, \dots, e_0]))}{\tilde{M}_m(K \times_m [e_m, \dots, e_0])} 1_{m[e_m, \dots, e_0]},$$

and ϕ is the modulus of uniform continuity of $p_e|_{K_{i(e)}}$. Then

$$\int X_m dM = \tilde{M}_m(Q_m) \leq a^{\frac{-m+1}{2}}.$$

Set

$$\Omega_m := \left\{ \sigma \in \Sigma : X_m(\sigma) > a^{\frac{-m+1}{4}} \right\}.$$

Then

$$a^{\frac{-m+1}{4}} M(\Omega_m) \leq \int_{\Omega_m} X_m dM \leq a^{\frac{-m+1}{2}}.$$

Hence

$$M(\Omega_m) \leq a^{\frac{-m+1}{4}}.$$

Set

$$\Omega := \bigcap_{n \leq 0} \bigcup_{m \leq n} \Omega_m.$$

Then $X_m(\sigma) \rightarrow 0$ for all $\sigma \in \Sigma \setminus \Omega$, and

$$M(\Omega) = 0,$$

by the Borel-Cantelli argument. Since each $p_e|_{K_{i(e)}}$ is uniformly continuous and $M(Y) = 1$ implies that $Y_m \rightarrow F$ M -a.e., we conclude, by (4), that

$$|E_M(1_{[e]}|\mathcal{F}_m) - p_e \circ F| \rightarrow 0 \quad M\text{-a.e.}$$

□

Proposition 1 *Suppose $\mathcal{M}$ is a CMS with an invariant Borel probability measure μ such that $\sum_{i=1}^N \int_{K_i} d(x, x_i) d\mu(x) < \infty$ for some $x_i \in K$, $i = 1, \dots, N$, and $M(Y) = 1$, where M is the generalized Markov measure associated with $\mathcal{M}$ and μ . Then the following hold.*

- (i) M is an equilibrium state for u .
- (ii) $P(u) = 0$.

Proof. By Lemma 6,

$$E_M(1_{1[e]}|\mathcal{F}) = p_e \circ F \text{ } M\text{-a.e. for all } e \in E.$$

For each $e \in E$, set

$$\dot{g}_{Me}(\sigma) := \begin{cases} p_e \circ F(\sigma) & \text{if } \sigma \in Y \\ 0 & \text{if } \sigma \in \Sigma \setminus Y. \end{cases}$$

Since $M(Y) = 1$, each $\dot{g}_{Me}$ is a version of g_{Me} which satisfies the hypothesis of Lemma 5 and $u = \log \dot{g}_M$, where $\dot{g}_M(\sigma) := \dot{g}_{M\sigma_1}(\sigma)$ for all $\sigma \in \Sigma$. Hence, by Lemma 5, M is an equilibrium state for u , and also holds (ii). $\square$

Proposition 2 *Let $\Lambda \in P_S(\Sigma)$ be an equilibrium state for u . Then*

$$U^*F(\Lambda) = F(\Lambda).$$

Proof. Since Λ is an equilibrium state for u , $\Lambda(Y) = 1$ (otherwise $h_\Lambda(S) + \Lambda(u) = -\infty < 0$). Furthermore, by Lemma 5, $g_\Lambda = \exp u$ Λ -a.e.. Hence, we can assume without loss of generality that $g_\Lambda(\sigma) = \exp u(\sigma)$ for all $\sigma \in Y$.

Now, let $f \in C_B(K)$ and $_{-1}[e_0, e] \cap Y \neq \emptyset$. Then, by the shift invariance of Λ and the pull-out property of the conditional expectation,

$$\begin{aligned} \int_{_{-1}[e_0, e]} f \circ F \, d\Lambda &= \int_{_0[e_0]} 1_{1[e]} f \circ F \circ S \, d\Lambda = \int_{_0[e_0]} 1_{1[e]} f \circ w_e \circ F \, d\Lambda \\ &= \int_{_0[e_0]} g_{\Lambda e} f \circ w_e \circ F \, d\Lambda. \end{aligned}$$

Let $\sigma \in _0[e_0] \cap Y$. Since $_0[e_0, e] \cap Y \neq \emptyset$, there exists $\sigma' \in Y$ such that $\sigma' = (\dots, \sigma_{-2}, \sigma_{-1}, e_0, e, \sigma'_2, \sigma'_3, \dots)$. Hence $g_{\Lambda e}(\sigma) = g_\Lambda(\sigma') = \exp u(\sigma') = p_e \circ F(\sigma)$. Thus

$$g_{\Lambda e}(\sigma) = p_e \circ F(\sigma) \text{ for all } \sigma \in _0[e_0] \cap Y.$$

Therefore

$$\int_{_0[e_0]} g_{\Lambda e} f \circ w_e \circ F \, d\Lambda = \int_{_0[e_0]} p_e \circ F f \circ w_e \circ F \, d\Lambda.$$

Summing for all $e_0 \in E$ gives

$$\int_{_0[e]} f \circ F \, d\Lambda = \int_{_0[e]} p_e \circ F f \circ w_e \circ F \, d\Lambda.$$

Hence,

$$F(\Lambda)(f) = \int f \circ F \, d\Lambda = \sum_{e \in E} \int p_e f \circ w_e \, dF(\Lambda) = U^*F(\Lambda)(f)$$

as desired. $\square$

Theorem 2 *Suppose CMS $\mathcal{M}$ has a unique invariant Borel probability measure μ and $M(Y) = 1$, where M is the associated generalized Markov measure. Then the following hold.*

- (i) *M is a unique equilibrium state for the energy function u ,*
- (ii) *$F(M) = \mu$,*
- (iii) *$h_M(S) = -\sum_{e \in E} \int_{K_{i(e)}} p_e \log p_e \, d\mu$.*

Proof. By Theorem 1 (iv), $\sum_{i=1}^N \int_{K_i} d(x, x_i) d\mu(x) < \infty$ for all $x_i \in K_i$, $i = 1, \dots, N$. Therefore, by Lemma 6,

$$E_M(1_{[e]} | \mathcal{F}) = p_e \circ F \text{ } M\text{-a.e. for all } e \in E.$$

Hence, by Proposition 1, M is an equilibrium state for u .

Now, suppose Λ is another equilibrium state for u . This implies that $\Lambda(Y) = 1$ (otherwise $h_\Lambda(S) + \Lambda(u) = -\infty < 0$). Also, by Lemma 5, $g_\Lambda(\sigma) = \dot{g}_M(\sigma)$ for Λ -a.e. $\sigma \in \Sigma$. Hence, we can assume, without loss of generality, that

$$g_{\Lambda\sigma_1}(\sigma) = p_{\sigma_1} \circ F(\sigma) \text{ for all } \sigma \in Y.$$

Let $[e_1, \dots, e_n] \subset \Sigma$ be a cylinder set such that $[e_1, \dots, e_n] \cap Y \neq \emptyset$. By the shift-invariance of Λ ,

$$\Lambda([e_1, \dots, e_n]) = \int_{-n+2[e_1, \dots, e_{n-1}]} 1_{[e_n]} d\Lambda = \int_{-n+2[e_1, \dots, e_{n-1}]} g_{\Lambda e_n} d\Lambda.$$

Let $\sigma \in -n+2[e_1, \dots, e_{n-1}] \cap Y$. Then there exists $\sigma' \in Y$ such that $\sigma' = (\dots, \sigma_{-1}, \sigma_0, e_n, \sigma'_2, \sigma'_3, \dots)$. Hence, $g_{\Lambda e_n}(\sigma) = g_{\Lambda\sigma'_1}(\sigma') = p_{\sigma'_1} \circ F(\sigma') = p_{e_n} \circ F(\sigma)$. We conclude that

$$g_{\Lambda e_n}(\sigma) = p_{e_n} \circ F(\sigma) \text{ for all } \sigma \in -n+2[e_1, \dots, e_{n-1}] \cap Y.$$

Note that F is $\mathcal{F}$ -measurable and $F(S\sigma) = w_{\sigma_1}(F(\sigma))$ for all $\sigma \in Y$. Therefore,

$$\begin{aligned} & \int_{-n+2[e_1, \dots, e_{n-1}]} g_{\Lambda e_n} d\Lambda \\ &= \int_{-n+2[e_1, \dots, e_{n-1}]} p_{e_n} \circ F d\Lambda \\ &= \int_{-n+3[e_1, \dots, e_{n-2}]} 1_{[e_{n-1}]} p_{e_n} \circ F \circ S d\Lambda \\ &= \int_{-n+3[e_1, \dots, e_{n-2}]} 1_{[e_{n-1}]} p_{e_n} \circ w_{e_{n-1}} \circ F d\Lambda. \end{aligned}$$

By using the pull-out property of the conditional expectation and repeating the above argumentation, we obtain that

$$\begin{aligned}
& \int_{-n+3[e_1, \dots, e_{n-2}]} 1_{[e_{n-1}]} p_{e_n} \circ w_{e_{n-1}} \circ F d\Lambda \\
&= \int_{-n+3[e_1, \dots, e_{n-2}]} p_{e_{n-1}} \circ F p_{e_n} \circ w_{e_{n-1}} \circ F d\Lambda \\
&\quad \cdot \\
&\quad \cdot \\
&\quad \cdot \\
&= \int p_{e_1} \circ F p_{e_2} \circ w_{e_1} \circ F \dots p_{e_n} \circ w_{e_{n-1}} \circ \dots \circ w_{e_1} \circ F d\Lambda \\
&= \int p_{e_1} p_{e_2} \circ w_{e_1} \dots p_{e_n} \circ w_{e_{n-1}} \circ \dots \circ w_{e_1} dF(\Lambda).
\end{aligned}$$

Thus, the equality $\Lambda = M$ will follow from $F(\Lambda) = \mu$, but this follows by the uniqueness of the invariant measure μ , since $U^*F(\Lambda) = F(\Lambda)$ by Proposition 2. Thus, the claims (i) and (ii) hold true. By Proposition 1 (ii) and Lemma 6,

$$\begin{aligned}
h_M(S) &= - \sum_{e \in E} \int 1_{[e]} \log p_e \circ F dM \\
&= - \sum_{e \in E} \int p_e \circ F \log p_e \circ F dM \\
&= - \sum_{e \in E} \int_{K_{i(e)}} p_e \log p_e d\mu.
\end{aligned}$$

This proves (iii). $\square$

Corollary 1 *Suppose $(K_{i(e)}, w_e, p_e)_{e \in E}$ is an irreducible CMS such that each $p_e|_{K_{i(e)}}$ is Dini-continuous and bounded away from zero. Then the following hold.*

- (i) *The generalized Markov measure M is a unique equilibrium state for the energy function u ,*
- (ii) *$P(u) = 0$,*
- (iii) *$F(M) = \mu$,*
- (iv) *$h_M(S) = - \sum_{e \in E} \int_{K_{i(e)}} p_e \log p_e d\mu$.*

Proof. By Theorem 2 in [13], the CMS has a unique invariant Borel probability measure. Since $M(Y) = 1$ by Lemma 3 (ii), the claims follow by Theorem 2. $\square$

Remark 3 The author would like to point out that a similar entropy formula as that proved in Theorem 2 (iv) plays a central role in the recent book of Wojciech Ślomeczynski [10].

Finally, we would like to make some remarks on why the result presented here might be interesting for the general theory of thermodynamic formalism.

Remark 4 First of all, recall that the theory of equilibrium states is presented usually only for upper semicontinuous energy functions (see e.g. [6]). The uniqueness of an equilibrium state is known on sub-shifts of finite type in general only for energy functions satisfying some stronger continuity conditions, e.g. Dini-continuity (note that the Dini-continuity and the *regularity* of a function coincide on one-dimensional lattices) [6], [9].

If all maps w_e of $\mathcal{M}$ are contractive and all probabilities $p_e|_{K_{i(e)}}$ are Dini-continuous, then the coding map F is defined everywhere on Σ_G and is Hölder-continuous. Hence, the energy function u is upper-semicontinuous and $u|_{\Sigma_G}$ is Dini-continuous (easy to check, since $\log x \leq x - 1$). In this case, Corollary 1 (i) fits nicely into the well known thermodynamic formalism.

Now, let us consider u if w_e 's are contractive only on average. In this case, Y is not necessarily closed (see Example 3), i.e. u is not necessarily upper semicontinuous. Therefore, even the existence of an equilibrium state for u is not guaranteed by the existing thermodynamic formalism. Moreover, by Lemma 2, we only know that for all $\epsilon > 0$ there exists $Q \subset Y$ with $M(Q) > 1 - \epsilon$ such that $u|_Q$ is Dini-continuous, but the sum $\sum_{k=1}^{\infty} \phi_Q(2^{-k})$, where ϕ_Q is the modulus of uniform continuity of $u|_Q$, increases if we choose Q larger.

Summing up, Corollary 1 shows that the general contractive Markov systems considered here still inherit some of their thermodynamic properties from finite Markov chains, even though their energy function belongs to a class which, as far as the author is aware, is not considered by the existing theory of thermodynamic formalism.

Remark 5 An important result of the thermodynamic formalism on topologically mixing subshifts of finite type is that for a Dini-continuous energy function the unique equilibrium state can be obtained as a unique *Gibbs state* for the same energy function (see e.g. [6], [9]).

If all w_e 's are contractive, the energy function $u|_{\Sigma_G}$ is Dini-continuous (as in Remark 4) and therefore, by Corollary 1 (i), M is also a unique Gibbs state for u . However, Y is not necessarily a subshift of finite type (not necessarily closed) and $u|_Y$ is not necessarily Dini-continuous if the maps are contractive only on average. We do not know in this case whether the measure M still can be constructed as a unique Gibbs state.

Acknowledgements

I would like to thank: EPSRC and School of Mathematics and Statistics of University of St Andrews for providing me with a scholarship and excellent working conditions in St Andrews, the anonymous referees for suggestions on

improvements for this paper. Also, I would like to thank Barry Ridge and Wang Yang for their help in the production of this paper.

References

- [1] M. F. Barnsley, S. G. Demko, J. H. Elton and J. S. Geronimo, Invariant measure for Markov processes arising from iterated function systems with place-dependent probabilities, *Ann. Inst. Henri Poincaré* **24** (1988) 367-394.
- [2] R. M. Dudley, *Real analysis and probability*, The Warsworth & Brooks/Cole Mathematics Series, California (1989).
- [3] J. H. Elton, An ergodic theorem for iterated maps, *Ergod. Th. & Dynam. Sys.* **7** (1987) 481-488.
- [4] R. Isaac, Markov processes and unique stationary probability measures, *Pacific J. Math.* **12** (1962) 273-286.
- [5] M. Keane, Strongly Mixing g -Measures, *Inventiones math.* **16** (1972), 309-324.
- [6] G. Keller, *Equilibrium States in Ergodic Theory*, Student Texts **42** LMS, Cambridge University Press (1998).
- [7] F. Ledrappier, Principe variationnel et systèmes dynamiques symboliques, *Z. Wahrscheinlichkeitstheorie verw. Gebiete* **30** (1974) 185-202.
- [8] R. D. Mauldin and S. C. Williams, Hausdorff dimension in graph directed constructions, *Trans. AMS* **309** (1988) 811-829.
- [9] D. Ruelle, *Thermodynamic formalism. The mathematical structures of classical equilibrium statistical mechanics*. Encyclopedia of Mathematics and its Applications, 5. Addison-Wesley Publishing Co., Reading, Mass.(1978).
- [10] W. Ślomczyński, *Dynamical entropy, Markov operators, and iterated function systems*, Rozprawy Habilitacyjne Uniwersytetu Jagiellońskiego Nr 362, Wydawnictwo Uniwersytetu Jagiellońskiego (2003).
- [11] P. Walters, Ruelle's Operator Theorem and g -measures, *Trans. Am. Math. Soc.* **214** (1975), 375-387.
- [12] P. Walters, *An introduction to Ergodic Theory*, Springer-Verlag (1982).
- [13] I. Werner, Contractive Markov systems, *J. London Math. Soc.* **71** (2005), no. 1, 236-258.
- [14] I. Werner, Contractive Markov system with constant Probabilities, *J. Theoret. Prob.* **18** (2005), no. 2, 469-479.

- [15] I. Werner, Coding map for a contractive Markov system, *Math. Proc. Camb. Phil. Soc.* to appear 140 (2) (2006).
- [16] I. Werner, Ergodic theorem for a contractive Markov system, *Nonlinearity* **17** (2004) 2303-2313.